

## Vascular ADAM17 (a Disintegrin and Metalloproteinase Domain 17) Is Required for Angiotensin II/ $\beta$ -Aminopropionitrile–Induced Abdominal Aortic Aneurysm

Tatsuo Kawai,\* Takehiko Takayanagi,\* Steven J. Forrester, Kyle J. Preston, Takashi Obama, Toshiyuki Tsuji, Tomonori Kobayashi, Michael J. Boyer, Hannah A. Cooper, Hang Fai Kwok, Tomoki Hashimoto, Rosario Scalia, Victor Rizzo, Satoru Eguchi

**Abstract**—Angiotensin II (AngII)–activated epidermal growth factor receptor has been implicated in abdominal aortic aneurysm (AAA) development. In vascular smooth muscle cells (VSMCs), AngII activates epidermal growth factor receptor via a metalloproteinase, ADAM17 (a disintegrin and metalloproteinase domain 17). We hypothesized that AngII-dependent AAA development would be prevented in mice lacking ADAM17 in VSMCs. To test this concept, control and VSMC ADAM17-deficient mice were cotreated with AngII and a lysyl oxidase inhibitor,  $\beta$ -aminopropionitrile, to induce AAA. We found that 52.4% of control mice did not survive because of aortic rupture. All other surviving control mice developed AAA and demonstrated enhanced expression of ADAM17 in the AAA lesions. In contrast, all AngII and  $\beta$ -aminopropionitrile-treated VSMC ADAM17-deficient mice survived and showed reduction in external/internal diameters (51%/28%, respectively). VSMC ADAM17 deficiency was associated with lack of epidermal growth factor receptor activation, interleukin-6 induction, endoplasmic reticulum/oxidative stress, and matrix deposition in the abdominal aorta of treated mice. However, both VSMC ADAM17-deficient and control mice treated with AngII and  $\beta$ -aminopropionitrile developed comparable levels of hypertension. Treatment of C57Bl/6 mice with an ADAM17 inhibitory antibody but not with control IgG also prevented AAA development. In conclusion, VSMC ADAM17 silencing or systemic ADAM17 inhibition seems to protect mice from AAA formation. The mechanism seems to involve suppression of epidermal growth factor receptor activation. (*Hypertension*. 2017;70:959-963. DOI: 10.1161/HYPERTENSIONAHA.117.09822.) • [Online Data Supplement](#)

**Key Words:** aneurysm ■ angiotensin II ■ hypertension ■ rupture ■ signal transduction

Abdominal aortic aneurysm (AAA) is a significant cause of death in the elderly, whereas effective pharmacological intervention is not yet available.<sup>1</sup> Activation of the renin angiotensin II (AngII) system has been implicated in AAA development in human and animal models of AAA.<sup>2,3</sup> While the detailed molecular mechanism by which AngII promotes AAA development remains unclear, it seems to involve multiple cell types (leukocytes, vascular smooth muscle cells [VSMCs], and endothelial cells) and signaling responses, such as oxidative stress, induction of inflammatory cytokines, and activation of matrix metalloproteinases.<sup>4,5</sup> However, limited information is available regarding the cell type–specific mechanism critical for AAA development. Therefore, the aims of this study are to define an essential signaling mechanism for development of AAA in an AngII-dependent mouse model and to provide a novel therapeutic target in AAA.

We have demonstrated that a metalloproteinase, ADAM17 (a disintegrin and metalloproteinase domain 17), is required for AngII-induced epidermal growth factor receptor (EGFR) transactivation in VSMCs<sup>6</sup> and that the ADAM17/EGFR activation mediates vascular remodeling in mice infused with AngII.<sup>7,8</sup> Moreover, ADAM17 expression is enhanced in human AAA,<sup>9,10</sup> and ADAM17-silenced mice do not develop CaCl<sub>2</sub>-induced AAAs.<sup>9</sup>

$\beta$ -Aminopropionitrile (BAPN) is an inhibitor for lysyl oxidase, which cross-links elastin fibers and collagen fibers and plays a critical role in maintaining homeostasis of the elastic lamina.<sup>11</sup> While BAPN treatment alone does not promote AAA, it causes AAA development and rupture when combined with AngII via degeneration of elastic lamina.<sup>12</sup> In VSMCs, both ADAM17 and EGFR colocalize at caveolae, and AAA formation induced by AngII plus BAPN was attenuated

Received June 2, 2017; first decision June 22, 2017; revision accepted August 31, 2017.

From the Cardiovascular Research Center, Department of Physiology, Lewis Katz School of Medicine at Temple University, Philadelphia, PA (T. Kawai, T. Takayanagi, S.J.F., K.J.P., T.O., T. Tsuji, T. Kobayashi, M.J.B., H.A.C., R.S., V.R., S.E.); Faculty of Health Sciences, Macau Special Administrative Region, University of Macau, Taipa (H.F.K.); and Department of Anesthesia and Perioperative Care, University of California, San Francisco (T.H.).

\*These authors contributed equally to this work.

The online-only Data Supplement is available with this article at <http://hyper.ahajournals.org/lookup/suppl/doi:10.1161/HYPERTENSIONAHA.117.09822/-DC1>.

Correspondence to Satoru Eguchi, Cardiovascular Research Center, Lewis Katz School of Medicine at Temple University, 3500 N Broad St, Philadelphia, PA 19140, E-mail [seguchi@temple.edu](mailto:seguchi@temple.edu) or Victor Rizzo, Cardiovascular Research Center, Lewis Katz School of Medicine at Temple University, 3500 N Broad St, Philadelphia, PA 19140, E-mail [rizzov@temple.edu](mailto:rizzov@temple.edu)

© 2017 American Heart Association, Inc.

*Hypertension* is available at <http://hyper.ahajournals.org>

DOI: 10.1161/HYPERTENSIONAHA.117.09822

in caveolin-1-deficient mice.<sup>13</sup> Moreover, we have recently reported that pharmacological inhibition of EGFR prevents AAA development induced by AngII plus BAPN, which was associated with suppression of endoplasmic reticulum (ER) stress, oxidative stress, and interleukin-6 and matrix metalloproteinases-2 expression.<sup>10</sup> However, whether VSMC-specific ADAM17 expression is essential for signal transduction leading to AAA development or any pharmacological ADAM17 intervention prevents AAA development remains unclear. In the present study, we tested the hypothesis that genetic silencing of VSMC ADAM17 or systemic ADAM17 inhibitory antibody treatment prevents AAA formation induced by AngII plus BAPN.

## Materials and Methods

### Animal Protocol

All animal procedures were performed with prior approval from Temple University Institutional Animal Care and Use Committee and in accordance with National Institutes of Health Guide for the Care and Use of Laboratory Animals. Mice deficient in VSMC ADAM17 (ADAM17<sup>fllox/fllox</sup> Sm22 $\alpha$ -Cre<sup>+/-</sup>) were generated as reported.<sup>14</sup> At 8 to 10 weeks of age, male ADAM17<sup>fllox/fllox</sup> Sm22 $\alpha$ -Cre<sup>+/-</sup> mice and littermate male control, ADAM17<sup>fllox/fllox</sup> Sm22 $\alpha$ -Cre<sup>-/-</sup>, mice were infused with AngII (Bachem; 1  $\mu$ g/kg-per minute) for 4 weeks and BAPN (Tokyo Chemical Industry, 150 mg/kg-per day) for the first 2 weeks or control saline for 4 weeks (Alzet, Durect Corp).<sup>13</sup> In addition, 8- to 10-week-old male C57Bl6 mice (Jackson) were infused with AngII (1  $\mu$ g/kg per min) for 4 weeks and received BAPN in drinking water (1 mg/mL) for the first 2 weeks together with human/mouse cross-reactive ADAM17 inhibitory antibody A9B8<sup>15</sup> or control human IgG2 (Athens Research & Technology), which was solubilized in PBS and administered at 10 mg/kg-per day via intraperitoneal injection, at days 1, 7, 14, and 21. Control C57Bl6 mice were sham-operated.

The AngII plus BAPN-induced mouse model of aortic aneurysm reproducibly induces AAA with morphological and histological characteristics similar to human AAA, but without enhancing atherosclerosis, as seen in other AngII-dependent AAA models.<sup>12</sup> Aortic luminal diameter at maximal dilation was measured using high-resolution 2-dimensional imaging (B mode) with high-frequency ultrasound (VisualSonics Vevo2100) on day 0, 14, 21, and 28 of the study. The treatment protocols were blinded to the evaluator. Despite the original article reporting incidence of thoracic aortic aneurysm in this model (38%),<sup>12</sup> we consistently observed much less thoracic aortic aneurysm (0%–10%).<sup>10,13</sup> Therefore, quantitative evaluation was not performed in thoracic aortas. Blood pressure and heart rate were evaluated in the conscious state at day 28 by telemetry (DSI equipped with ADInstrument 6 software) via carotid catheter (PA-C10 transmitter). For animals that died before the completion of the study, necropsy was performed when possible to define the causes of death. The thoracic aorta or abdominal aorta rupture as a cause of death was defined with the presence of hematoma in thoracic or abdominal cavity, respectively. After 28 days, mice were euthanized, perfused with formalin, and dissected for tissue samples. Abdominal aortas were extracted and subjected to paraffin embedding. Sections were stained with a standard Masson's trichrome protocol<sup>8</sup> to distinguish medial area from adventitia. Images were visualized on an Olympus IX81 inverted microscope using an Olympus SC30 high-resolution camera and acquired with Olympus cellSens Entry 1.11 software.

### Immunohistochemistry

Sections from abdominal aortas were deparaffinized and blocked in 5% goat serum and 1% BSA for 1 hour at room temperature, incubated with primary antibody in PBS containing 1% BSA and 0.1% Tween 20 overnight at 4°C followed by biotinylated secondary antibody for 90 minutes at room temperature. The sources and dilutions of the primary antibodies used in this study are provided in Table S1 in the online-only Data Supplement. Slides were incubated with


avidin–biotin peroxidase complex for 30 minutes at room temperature, and staining was visualized with the substrate diaminobenzidine (Vector), which produced a brown color, and counterstained with hematoxylin. An equal concentration of control IgG was used side-by-side with each antibody to ensure staining specificity. Quantification of the antibody staining was performed as reported previously with subtraction of IgG background staining.<sup>13</sup> All images were visualized on Olympus SC30 high-resolution camera and acquired with Olympus cellSens Entry 1.11 software using the same exposure time. Images were loaded into the ImageJ program (<http://rsb.info.nih.gov/ij/>) for analysis. A region of interest was drawn around the entire aorta with the freehand selection tool. Adventitia was excluded from the quantification because the adventitia areas were limited in aortas except those with AAA. All images were set to the same hue, saturation, and brightness. The area and intensity (integrated density) in the region of interest were then measured and analyzed. Data were obtained from 3 to 4 nonoverlapping fields per aortic cross-section for each antibody (n=4 aortas per treatment or genotype). Results are presented as fold increase over control, which was set at 1.

### Statistical Analysis

Data from the groups were analyzed by 1-way analysis of variance with Tukey's multiple comparison test (C57Bl6 mice with basal, AAA treatment, or AAA treatment plus ADAM17 antibody), 2-way analysis of variance with Bonferroni post-tests (ADAM17 wild-type or deficient mice with or without AAA treatment), or log-rank (Mantel–Cox) test (Kaplan–Meier survival curves) using GraphPad Prism version 5.0C for Macintosh. Data were presented as mean $\pm$ SEM. Statistical significance was taken at  $P<0.05$ .

## Results

There was a significant difference in survival rates between mice deficient with VSMC ADAM17 (100%) and littermate control mice (47.6%) when treated with AngII plus BAPN over the 28-day observational period (Figure 1A). Based on necropsy, all confirmed deaths were caused by aortic rupture. Surviving control mice, as well as mice deficient in VSMC ADAM17, treated with AngII plus BAPN developed comparable degree of hypertension (Figure 1B; Table S2). All surviving AngII plus BAPN-treated control mice developed AAA, with significant enlargements of the maximum diameter of abdominal aortas. In contrast, VSMC ADAM17-deficient



**Figure 1.** **A**, Vascular smooth muscle cells (VSMC) ADAM17 (a disintegrin and metalloproteinase domain 17)-deficient mice survived from aortic rupture. Eight-wk-old VSMC ADAM17-deficient mice (ADAM17<sup>fllox/fllox</sup> sm22 $\alpha$ -Cre<sup>+/-</sup>, A17<sup>ff</sup>Cre<sup>+/-</sup>) and control mice (ADAM17<sup>fllox/fllox</sup> sm22 $\alpha$ -Cre<sup>-/-</sup>, A17<sup>ff</sup>Cre<sup>-/-</sup>) were infused with angiotensin II (1  $\mu$ g/kg-per minute, 4 wk) and  $\beta$ -aminopropionitrile (BAPN; 150 mg/kg-per day, first 2 wk) or saline for 4 wk. Percentage survival curve is shown (n=21). **B**, Telemetry recording of mean arterial blood pressure (MAP) on 4-wk infusion (n=5).

mice with AngII plus BAPN treatment had significantly less aortic diameter enhancement compared with control mice (Figure 2; Figures S1 and S2).

Histological analysis demonstrated that AAAs induced by AngII plus BAPN in control mice were associated with vascular fibrosis/matrix deposition and disruption of medial layer structures. In addition, enhanced EGFR activation, increased expression of ADAM17, matrix metalloproteinases-2, and interleukin-6, and enhanced ER stress (Lys-Asp-Glu-Leu), oxidative stress (nitro-tyrosine), and leukocyte infiltration (CD45) were observed with semiquantitative immunohistochemical staining. These AAA-associated responses were attenuated in VSMC ADAM17-deficient mice. However, tumor necrosis factor- $\alpha$  (TNF $\alpha$ ) expression did not show any statistical differences among the groups (Figures S3 through S6).


To ascertain that ADAM17 represents a novel therapeutic target to prevent AAA development, C57Bl6 mice treated with AngII plus BAPN were injected with a human/mouse cross-reactive ADAM17 inhibitory antibody. Distinct from VSMC ADAM17 silencing, there was no statistical difference

in the survival rate with the ADAM17 antibody treatment (Figure 3A). The ADAM17 antibody treatment did not alter hypertension induced by AngII plus BAPN (Figure 3B; Table S3). However, AAA development was significantly reduced in mice treated with the ADAM17 antibody compared with the control IgG treatment (Figure 4; Figures S7 through S9).


Discussion

Enhanced ADAM17 expression<sup>9</sup> and downstream EGFR activation<sup>10</sup> have been reported in human AAA samples. Requirement of ADAM17 in CaCl<sub>2</sub>-induced AAA has also been reported with inducible systemic ADAM17 deletion.<sup>9</sup> However, our study provides new information that AngII-dependent AAA development and rupture are markedly prevented in mice lacking VSMC ADAM17 and that pharmacological intervention of ADAM17 can attenuate AAA in a mouse model. Limited VSMC-specific mechanisms are known to contribute to AAA in animal models, which includes decreased catalase<sup>16</sup> and activation of Notch1,<sup>17</sup> whereas induction of hypoxia-inducible factor-1 $\alpha$  in VSMCs seems protective.<sup>18</sup>


In VSMCs stimulated with AngII, ADAM17-dependent shedding produces EGFR ligands, such as heparin-binding epidermal growth factor-like growth factor leading to EGFR transactivation.<sup>19</sup> Because EGFR inhibition also prevents AngII plus BAPN-mediated AAA development and rupture,<sup>10</sup> VSMC EGFR most likely mediates the ADAM17-dependent function in this mouse model of AAA. On activation, EGFR mediates several downstream responses in VSMCs, including oxidative stress,<sup>20</sup> ER stress,<sup>8</sup> and interleukin-6 induction.<sup>21</sup> These downstream effects were also evident in the present study, thus, likely contributing to AAA development (Figure S10).<sup>16,22,23</sup> In addition, enhanced vascular ER and oxidative stress further promote immune cell infiltration, which is also critical for AAA development.<sup>4</sup> Therefore, the findings demonstrated in this article suggest that all of these mechanisms are potentially mediated by a single but multifunctional metalloprotease ADAM17 specifically expressed in VSMCs.



**Figure 2.** Vascular smooth muscle cells (VSMC) ADAM17 (a disintegrin and metalloproteinase domain 17)-deficient mice did not develop angiotensin II (AngII)-dependent abdominal aortic aneurysm (AAA). VSMC ADAM17-deficient mice (A17<sup>ff</sup>sm22 $\alpha$ Cre<sup>+/-</sup>) and control littermate mice (Cre<sup>-/-</sup>) were infused with AngII plus  $\beta$ -aminopropionitrile (BAPN) or saline as in Figure 1. **A**, Measurements of maximal external width of abdominal aorta (AA) on fixation at 4 wk (Cre<sup>-/-</sup>, 2.29 $\pm$ 0.50 mm vs Cre<sup>+/-</sup>, 1.12 $\pm$ 0.23 mm with AngII plus BAPN; n=6). **B**, Weekly ultrasound evaluation of maximal abdominal aorta luminal diameter. The panels shown were at 4 wk. \*, †P<0.05 compared with control saline or AngII/BAPN infusion, respectively (each n=5). The measurements were blinded and performed by 2 evaluators confirming the reproducibility.



**Figure 3. A**, Survival data of C57Bl6 mice with ADAM17 (a disintegrin and metalloproteinase domain 17) antibody treatment. Eight-week-old C57Bl6 mice infused with angiotensin II (AngII; 1  $\mu$ g/kg-per min, 4 wk) and received  $\beta$ -aminopropionitrile (BAPN; 1 mg/mL in drinking water, first 2 wk) were intraperitoneally injected ADAM17 antibody A9B8 or control human IgG2 (10 mg/kg) at day 1, 7, 14, and 21. **B**, Telemetry recording of mean arterial blood pressure (MAP) on 4-wk infusion. Control C57Bl6 mice are sham-operated for minipump implantation (n=5).



**Figure 4.** ADAM17 (a disintegrin and metalloproteinase domain 17) antibody attenuated abdominal aortic aneurysm (AAA) development. C57Bl6 mice were received angiotensin II (AngII) plus  $\beta$ -aminopropionitrile (BAPN) together with ADAM17 antibody A9B8 or control IgG2 as in Figure 3. **A**, Measurements of maximal external width of abdominal aorta on fixation at 4 wk ( $n=6$ ). **B**, Weekly ultrasound evaluation of maximal abdominal aorta (AA) luminal diameter. \*,  $\dagger P < 0.05$  compared with control saline or AngII/BAPN infusion, respectively (each  $n=6-7$ ).

However, additional experiments are desired to track these mechanisms before the establishment of AAA. It is also ideal to use a distinct model of AAA to generalize our findings, which are limitations in this study.

While we confirmed a previously suggested EGFR-dependent mechanism as a downstream signal of VSMC ADAM17, our findings cannot exclude other and potentially new mechanism(s) through which ADAM17 contributes to AAA development. Other ADAM17 substrates, including TNF $\alpha$ , Notch1, and angiotensin-converting enzyme 2, may also participate in AAA pathology according to literature.<sup>17,24,25</sup> Note that while the protein expression analysis of TNF $\alpha$  did not show any enhancement in AAA, the experiment did not measure the potential conversion of pro-TNF $\alpha$  to mature and active TNF $\alpha$  by ADAM17. Moreover, ADAM17 has additional diverse substrates and many regulatory mechanisms.<sup>26</sup> Therefore, further research is required to identify potentially new pathways through which ADAM17 regulates AAA.

The AngII plus BAPN model consistently produces AAA associated with hypertension, but without enhancing atherosclerosis, as observed in other AngII-dependent AAA models, including hyperlipidemic mice.<sup>12</sup> Incidence of AAAs in normolipidemic mice with AngII infusion alone is low,<sup>27</sup> even though AngII infusion is sufficient for ADAM17 induction and EGFR activation in the vasculature, including the aorta.<sup>8</sup> Therefore, we surmise that ADAM17/EGFR activation is required to advance AAA but insufficient to initiate AAA, which requires an additional signal or condition such as those primed by BAPN or hyperlipidemia.

In the present study, a discrepancy was observed in survival rates between VSMC ADAM17 deletion with mixed background mice and systemic ADAM17 inhibition with C57BL6 mice. This may be because of significantly higher blood pressure in C57BL6 mice, regardless of the ADAM17 antibody or control IgG treatment. The discrepancy could also

be because of distinct genetic background of the mice and distinct cell type-specific roles (promoting versus preventing rupture in AAA) of ADAM17. In addition to the medial layer, enhanced ADAM17 expression was also observed in endothelium and adventitia of AAA, a finding that is in agreement with prior published work.<sup>13</sup> Because the promoter used to target ADAM17 is relatively specific to smooth muscle,<sup>14</sup> we assume that attenuation of AAA by silencing VSMC ADAM17 results in prevention of endothelial and adventitial ADAM17 induction by AngII plus BAPN. In addition to VSMC ADAM17, the ADAM17 antibody likely inhibited endothelial and adventitial ADAM17 in the present study. Therefore, the roles of endothelial and adventitial ADAM17 in AAA require further study.

## Perspectives

Our findings highlight the critical role of ADAM17 in mediating AAA development and rupture. We propose that VSMC ADAM17 is a needed component for EGFR transactivation contributing to ER stress and oxidative stress in AAA. Our results also indicate that ADAM17 inhibition can be a valuable treatment option for AAA. However, it remains to be explored how the VSMC ADAM17 signal communicates with other cell type-specific mechanisms presented before.

## Sources of Funding

This work was supported by National Institute of Health grants HL128324 (S. Eguchi and V. Rizzo), HL133248 (S. Eguchi), DK111042 (R. Scalia and S. Eguchi), F31HL127971 (S.J. Forrester), American Heart Association grants 16GRNT30410007 (S. Eguchi), 16GRNT30130013 (V. Rizzo), 15POST25550083 (K.J. Preston), and 16POST3051004 (T. Kawai), University of Macau grants SRG2014-0006-FHS (H.F. Kwok) and MYRG2015-00025-FHS (H.F. Kwok), and Science and Technology Development Fund of Macau FDCT 018/2015/A1 (H.F. Kwok).

## Disclosures

None.

## References

- Nordon IM, Hinchliffe RJ, Loftus IM, Thompson MM. Pathophysiology and epidemiology of abdominal aortic aneurysms. *Nat Rev Cardiol*. 2011;8:92-102. doi: 10.1038/nrcardio.2010.180.
- Golledge J, Norman PE. Pathophysiology of abdominal aortic aneurysm relevant to improvements in patients' management. *Curr Opin Cardiol*. 2009;24:532-538. doi: 10.1097/HCO.0b013e328330c2d3.
- Lu H, Rateri DL, Bruemmer D, Cassis LA, Daugherty A. Involvement of the renin-angiotensin system in abdominal and thoracic aortic aneurysms. *Clin Sci (Lond)*. 2012;123:531-543. doi: 10.1042/CS20120097.
- Davis FM, Rateri DL, Daugherty A. Abdominal aortic aneurysm: novel mechanisms and therapies. *Curr Opin Cardiol*. 2015;30:566-573. doi: 10.1097/HCO.0000000000000216.
- Emeto TI, Moxon JV, Au M, Golledge J. Oxidative stress and abdominal aortic aneurysm: potential treatment targets. *Clin Sci (Lond)*. 2016;130:301-315. doi: 10.1042/CS20150547.
- Ohtsu H, Dempsey PJ, Frank GD, Brailoiu E, Higuchi S, Suzuki H, Nakashima H, Eguchi K, Eguchi S. ADAM17 mediates epidermal growth factor receptor transactivation and vascular smooth muscle cell hypertrophy induced by angiotensin II. *Arterioscler Thromb Vasc Biol*. 2006;26:e133-e137. doi: 10.1161/01.ATV.0000236203.90331.d0.
- Takayanagi T, Forrester SJ, Kawai T, Obama T, Tsuji T, Elliott KJ, Nuti E, Rossello A, Kwok HF, Scalia R, Rizzo V, Eguchi S. Vascular ADAM17 as a novel therapeutic target in mediating cardiovascular hypertrophy and perivascular fibrosis induced by angiotensin II. *Hypertension*. 2016;68:949-955. doi: 10.1161/HYPERTENSIONAHA.116.07620.

8. Takayanagi T, Kawai T, Forrester SJ, Obama T, Tsuji T, Fukuda Y, Elliott KJ, Tilley DG, Davisson RL, Park JY, Eguchi S. Role of epidermal growth factor receptor and endoplasmic reticulum stress in vascular remodeling induced by angiotensin II. *Hypertension*. 2015;65:1349–1355. doi: 10.1161/HYPERTENSIONAHA.115.05344.
9. Kaneko H, Anzai T, Horiuchi K, Kohno T, Nagai T, Anzai A, Takahashi T, Sasaki A, Shimoda M, Maekawa Y, Shimizu H, Yoshikawa T, Okada Y, Yozu R, Fukuda K. Tumor necrosis factor- $\alpha$  converting enzyme is a key mediator of abdominal aortic aneurysm development. *Atherosclerosis*. 2011;218:470–478. doi: 10.1016/j.atherosclerosis.2011.06.008.
10. Obama T, Tsuji T, Kobayashi T, Fukuda Y, Takayanagi T, Taro Y, Kawai T, Forrester SJ, Elliott KJ, Choi ET, Daugherty A, Rizzo V, Eguchi S. Epidermal growth factor receptor inhibitor protects abdominal aortic aneurysm in a mouse model. *Clin Sci (Lond)*. 2014;128:559–565. doi: 10.1042/CS20140696.
11. Rodríguez C, Martínez-González J, Raposo B, Alcudia JF, Guadall A, Badimon L. Regulation of lysyl oxidase in vascular cells: lysyl oxidase as a new player in cardiovascular diseases. *Cardiovasc Res*. 2008;79:7–13. doi: 10.1093/cvr/cvn102.
12. Kanematsu Y, Kanematsu M, Kurihara C, Tsou TL, Nuki Y, Liang EI, Makino H, Hashimoto T. Pharmacologically induced thoracic and abdominal aortic aneurysms in mice. *Hypertension*. 2010;55:1267–1274. doi: 10.1161/HYPERTENSIONAHA.109.140558.
13. Takayanagi T, Crawford KJ, Kobayashi T, Obama T, Tsuji T, Elliott KJ, Hashimoto T, Rizzo V, Eguchi S. Caveolin 1 is critical for abdominal aortic aneurysm formation induced by angiotensin II and inhibition of lysyl oxidase. *Clin Sci (Lond)*. 2014;126:785–794. doi: 10.1042/CS20130660.
14. Weskamp G, Mendelson K, Swendeman S, Le Gall S, Ma Y, Lyman S, Hinoki A, Eguchi S, Guaiquil V, Horiuchi K, Blobel CP. Pathological neovascularization is reduced by inactivation of ADAM17 in endothelial cells but not in pericytes. *Circ Res*. 2010;106:932–940. doi: 10.1161/CIRCRESAHA.109.207415.
15. Kwok HF, Botkjaer KA, Tape CJ, Huang Y, McCafferty J, Murphy G. Development of a 'mouse and human cross-reactive' affinity-matured exosite inhibitory human antibody specific to TACE (ADAM17) for cancer immunotherapy. *Protein Eng Des Sel*. 2014;27:179–190. doi: 10.1093/protein/gzu010.
16. Parastatidis I, Weiss D, Joseph G, Taylor WR. Overexpression of catalase in vascular smooth muscle cells prevents the formation of abdominal aortic aneurysms. *Arterioscler Thromb Vasc Biol*. 2013;33:2389–2396. doi: 10.1161/ATVBAHA.113.302175.
17. Sachdeva J, Mahajan A, Cheng J, Baeten JT, Lilly B, Kuivaniemi H, Hans CP. Smooth muscle cell-specific Notch1 haploinsufficiency restricts the progression of abdominal aortic aneurysm by modulating CTGF expression. *PLoS One*. 2017;12:e0178538. doi: 10.1371/journal.pone.0178538.
18. Imanishi M, Chiba Y, Tomita N, Matsunaga S, Nakagawa T, Ueno M, Yamamoto K, Tamaki T, Tomita S. Hypoxia-inducible factor-1 $\alpha$  in smooth muscle cells protects aortic aneurysms-brief report. *Arterioscler Thromb Vasc Biol*. 2016;36:2158–2162. doi: 10.1161/ATVBAHA.116.307784.
19. Forrester SJ, Kawai T, O'Brien S, Thomas W, Harris RC, Eguchi S. Epidermal growth factor receptor transactivation: mechanisms, pathophysiology, and potential therapies in the cardiovascular system. *Annu Rev Pharmacol Toxicol*. 2016;56:627–653. doi: 10.1146/annurev-pharmtox-070115-095427.
20. Seshiah PN, Weber DS, Rocic P, Valppu L, Taniyama Y, Griendling KK. Angiotensin II stimulation of NAD(P)H oxidase activity: upstream mediators. *Circ Res*. 2002;91:406–413.
21. Ichiki T. Role of cAMP response element binding protein in cardiovascular remodeling: good, bad, or both? *Arterioscler Thromb Vasc Biol*. 2006;26:449–455. doi: 10.1161/01.ATV.0000196747.79349.d1.
22. Qin Y, Wang Y, Liu O, Jia L, Fang W, Du J, Wei Y. Tauroursodeoxycholic acid attenuates angiotensin II induced abdominal aortic aneurysm formation in apolipoprotein E-deficient mice by inhibiting endoplasmic reticulum stress. *Eur J Vasc Endovasc Surg*. 2017;53:337–345. doi: 10.1016/j.ejvs.2016.10.026.
23. Harrison SC, Smith AJ, Jones GT, et al; Aneurysm Consortium. Interleukin-6 receptor pathways in abdominal aortic aneurysm. *Eur Heart J*. 2013;34:3707–3716. doi: 10.1093/eurheartj/ehs354.
24. Xiong W, MacTaggart J, Knispel R, Worth J, Persidsky Y, Baxter BT. Blocking TNF- $\alpha$  attenuates aneurysm formation in a murine model. *J Immunol*. 2009;183:2741–2746. doi: 10.4049/jimmunol.0803164.
25. Thatcher SE, Zhang X, Howatt DA, Yiannikouris F, Gurley SB, Ennis T, Curci JA, Daugherty A, Cassis LA. Angiotensin-converting enzyme 2 decreases formation and severity of angiotensin II-induced abdominal aortic aneurysms. *Arterioscler Thromb Vasc Biol*. 2014;34:2617–2623. doi: 10.1161/ATVBAHA.114.304613.
26. Xu J, Mukerjee S, Silva-Alves CR, Carvalho-Galvão A, Cruz JC, Balarini CM, Braga VA, Lazartigues E, França-Silva MS. A disintegrin and metalloprotease 17 in the cardiovascular and central nervous systems. *Front Physiol*. 2016;7:469. doi: 10.3389/fphys.2016.00469.
27. Remus EW, O'Donnell RE Jr, Rafferty K, Weiss D, Joseph G, Csizsar K, Fong SF, Taylor WR. The role of lysyl oxidase family members in the stabilization of abdominal aortic aneurysms. *Am J Physiol Heart Circ Physiol*. 2012;303:H1067–H1075. doi: 10.1152/ajpheart.00217.2012.

## Novelty and Significance

### What Is New?

- Systemic and vascular smooth muscle cells-specific ADAM17 (a disintegrin and metalloproteinase domain 17) inhibition established a role for ADAM17 in angiotensin II-dependent abdominal aortic aneurysm (AAA) development independent of hypertension in mice.
- The concept of vascular ADAM17 in mediating the epidermal growth factor receptor pathway, oxidative stress, endoplasmic reticulum stress, and inflammation was presented.

### What Is Relevant?

- Results indicating prevention of AAA but not hypertension by ADAM17 inhibition provide a foundation to seek a potential therapy to prevent AAA development.

- The vascular dominant ADAM17 signal transduction highlights the importance of vascular signal transduction for AAA formation.

### Summary

In angiotensin II plus  $\beta$ -aminopropionitrile-treated vascular smooth muscle cells ADAM17-deficient mice, AAA development and rupture were prevented compared with treated control mice. These aortas with AAA showed vascular epidermal growth factor receptor activation and induction of ER stress and oxidative stress markers, which were attenuated in ADAM17-deficient mice. ADAM17 inhibitory antibody was used to confirm the contribution of ADAM17 in AAA pathology.