

# β2-Adrenergic receptor-dependent chemokine receptor 2 expression regulates leukocyte recruitment to the heart following acute injury

Laurel A. Grisanti<sup>a</sup>, Christopher J. Traynham<sup>a</sup>, Ashley A. Repas<sup>a</sup>, Erhe Gao<sup>a</sup>, Walter J. Koch<sup>a</sup>, and Douglas G. Tilley<sup>a,1</sup>

<sup>a</sup>Center for Translational Medicine, Lewis Katz School of Medicine, Temple University, Philadelphia, PA 19140

Edited by Brian K. Kobilka, Stanford University School of Medicine, Stanford, CA, and approved November 20, 2016 (received for review July 6, 2016)

Following cardiac injury, early immune cell responses are essential for initiating cardiac remodeling and tissue repair. We previously demonstrated the importance of β2-adrenergic receptors (β2ARs) in the regulation of immune cell localization following acute cardiac injury, with deficient leukocyte infiltration into the damaged heart. The purpose of this study was to investigate the mechanism by which immune cell-expressed β2ARs regulate leukocyte recruitment to the heart following acute cardiac injury. Chemokine receptor 2 (CCR2) expression and responsiveness to C-C motif chemokine ligand 2 (CCL2)-mediated migration were abolished in β2AR knock-out (KO) bone marrow (BM), both of which were rescued by β2AR reexpression. Chimeric mice lacking immune cell-specific CCR2 expression, as well as wild-type mice administered a CCR2 antagonist, recapitulated the loss of monocyte/macrophage and neutrophil recruitment to the heart following myocardial infarction (MI) observed in mice with immune cell-specific β2AR deletion. Conversely to β2AR ablation, β2AR stimulation increased CCR2 expression and migratory responsiveness to CCL2 in BM. Mechanistically, G protein-dependent β2AR signaling was dispensable for these effects, whereas β-arrestin2-biased β2AR signaling was required for the regulation of CCR2 expression. Additionally, activator protein 1 (AP-1) was shown to be essential in mediating CCR2 expression in response to β2AR stimulation in both murine BM and human monocytes. Finally, reconstitution of β2ARKO BM with rescued expression of a β-arrestin-biased β2AR in vivo restored BM CCR2 expression as well as cardiac leukocyte infiltration following MI. These results demonstrate the critical role of β-arrestin2/AP-1-dependent β2AR signaling in the regulation of CCR2 expression and recruitment of leukocytes to the heart following injury.

β2-adrenergic receptor | leukocyte | C-chemokine receptor 2 | cardiac injury | β-arrestin

Healing following ischemic cardiac injury is highly regulated by immune responses, with impairments or exacerbations in inflammation leading to alterations in infarct expansion, remodeling, and ultimately cardiac function (1). Cells of the innate immune system including monocytes/macrophages, mast cells, and neutrophils play critical roles in infarct healing through tissue phagocytosis and activation of reparative responses. Recruitment and trafficking of these leukocytes to the heart following acute injury occur through the action of chemokines on their receptors to promote their migration to the site of injury (2), and have been the focus of much research in recent years (1, 3).

Sympathetic activity is important for regulating immune responses, primarily through the β2-adrenergic receptor (β2AR) subtype (4–6). Recently, we showed that immune cell-expressed β2AR is required for leukocyte recruitment to the heart following acute myocardial infarction (MI), without which the heart cannot mount a repair response, ultimately undergoing rupture (7). Because chemokine receptors play a critical role in migration and infiltration of leukocyte populations, we hypothesized that immune cell-expressed chemokine receptor activity and/or expression may be altered in the absence of β2AR, thereby impairing leukocyte migration to the injured heart.

The impact of immune cell-specific β2AR expression on chemokine receptor expression and leukocyte infiltration following MI was investigated through the use of chimeric mice, wherein bone marrow transplant (BMT) recipient mice received bone marrow from β2ARKO donor mice. Through the use of these chimeric mice, we demonstrate that β2AR is critical in regulating chemokine receptor 2 (CCR2) expression, and responsiveness to its ligand C-C motif chemokine ligand 2 (CCL2), via a β-arrestin2 (βARR2)-biased signaling pathway involving activator protein 1 (AP-1). These results highlight the importance of β2AR in regulating immune cell expression of CCR2, thereby impacting the ability of leukocytes to respond to acute cardiac injury.

## Results

**CCR2 Expression and Migratory Responsiveness Are Abolished in β2ARKO BM.** We recently observed decreased leukocyte infiltration into the hearts of chimeric mice lacking immune cell-expressed β2AR following MI (7). Chemokines produced following injury are important for recruitment of immune cells, through their action on chemokine receptors. Thus, to assess whether differences in chemokine receptor expression could contribute to alterations in leukocyte infiltration in β2ARKO BMT mouse hearts post-MI, reverse transcription-quantitative PCR (RT-qPCR) was used to examine those known to play an important role in immune cell migration following acute cardiac injury (Table 1 and Table S1). β2ARKO BM had significantly decreased expression of CCR2 and C-X-C motif chemokine receptor 4 (CXCR4) compared with WT BM. To test the impact of

## Significance

The sympathetic nervous system influences various immune cell functions, in particular via β2-adrenergic receptor (β2AR) signaling. Although immune cell recruitment is critical for cardiac repair following ischemia, the impact of β2AR on this process is unclear. We describe how immune cell-specific β2AR depletion ablates chemokine receptor 2 (CCR2) expression and leukocyte recruitment to the heart postischemia. Reciprocally, β2AR activation increases CCR2 expression and responsiveness in a β-arrestin-dependent manner, and expression of a β-arrestin-biased β2AR in β2AR-depleted immune cells restores CCR2 levels and leukocyte recruitment to the postischemic heart. These results highlight the potential utility of next-generation β-arrestin-biased β2AR ligands to selectively modulate leukocyte responsiveness, and suggest that β-blockers, used commonly in peri/postischemic patients, may impact leukocyte-mediated repair mechanisms.

Author contributions: L.A.G. and D.G.T. designed research; L.A.G., A.A.R., and E.G. performed research; C.J.T. and W.J.K. contributed new reagents/analytic tools; L.A.G. analyzed data; and L.A.G. and D.G.T. wrote the paper.

Conflict of interest statement: W.J.K. and D.G.T. have equity in Renovacor, Inc., which has neither funded this study nor has a relevant product related to this study.

This article is a PNAS Direct Submission.

<sup>1</sup>To whom correspondence should be addressed. Email: douglas.tilley@temple.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1611023114/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1611023114/-DCSupplemental).

**Table 1. Effects of  $\beta$ 2ARKO on chemokine receptor expression**

Chemokine receptor	WT BMT	$\beta$ 2ARKO BMT
CCR1	1.00 $\pm$ 0.20	1.06 $\pm$ 0.13
CCR2	1.00 $\pm$ 0.21	0.27 $\pm$ 0.07*
CCR5	1.00 $\pm$ 0.16	1.32 $\pm$ 0.18
CXCR1	1.00 $\pm$ 0.54	1.71 $\pm$ 0.56
CXCR2	1.00 $\pm$ 0.51	1.63 $\pm$ 0.62
CXCR4	1.00 $\pm$ 0.06	0.32 $\pm$ 0.02*
CXCR7	1.00 $\pm$ 0.03	0.90 $\pm$ 0.04
CXC3CR1	1.00 $\pm$ 0.12	1.09 $\pm$ 0.14
CD45	1.00 $\pm$ 0.22	0.90 $\pm$ 0.30

RT-qPCR analysis of changes in expression of chemokine receptor transcripts in reconstituted WT or  $\beta$ 2ARKO BM from transplanted mice.  $n = 4-8$ .

\* $P < 0.001$  vs. WT BMT, two-tailed unpaired  $t$  test.


these altered chemokine receptor levels, we performed in vitro migration assays, wherein  $\beta$ 2ARKO BM displayed decreased migration toward CCL2 (MCP-1), the ligand for CCR2, with no difference in migratory responses to CCL3 or C-X-C motif chemokine ligand 12 (CXCL12), a CXCR4 ligand (Fig. 1*A* and *B*). Lentiviral-mediated restoration of  $\beta$ 2AR expression in  $\beta$ 2ARKO BM restored CCR2 expression to endogenous levels (Fig. 1*C*) as well as the migratory response to CCL2, without affecting migration to CCL3 or CXCL12 (Fig. 1*A* and *B*). CCR2 antagonism blocked migration toward CCL2 following  $\beta$ 2AR rescue but had no effect on CCL3 and CXCL12 responses, confirming that this response was CCR2-dependent.

**Specific Ablation of CCR2 Reduces Leukocyte Recruitment to the Heart Following MI.** Based on our in vitro assessment of the impact of  $\beta$ 2AR deletion on CCR2 expression and BM cell migration, we sought to determine whether CCR2 inhibition in vivo, either pharmacologically or genetically, could recapitulate the impaired leukocyte post-MI infiltration phenotype observed in  $\beta$ 2ARKO BMT mice (7). To assess this, WT mice underwent sham or MI surgery followed by daily injections with vehicle or CCR2 antagonist (2 mg·kg<sup>-1</sup>·d<sup>-1</sup>), or underwent irradiation and received WT,  $\beta$ 2ARKO, or CCR2KO BM 1 mo before surgery (Fig. S1). Analysis of infarct size 4 d post-MI showed no differences between groups, confirming similar surgical conditions for all groups of animals (Fig. S2). Immunohistochemistry was performed on heart sections 4 d postsurgery to quantify infiltration of immune cell populations in sham hearts and the remote (Fig. S3), border (Fig. 2 and Fig. S2*E-H*), and infarct (Fig. S4) zones of MI hearts. Both pharmacological CCR2 antagonism (Fig. S2*E* and *F*) and genetic CCR2 deletion (Fig. 2*A* and *B*) significantly reduced the infiltration of monocytes/macrophages (CD68<sup>+</sup> cells) and neutrophils [myeloperoxidase (MPO)<sup>+</sup> cells; Fig. 2*A* and *D* and Fig. S2*E* and *H*] to the border and infarct (Fig. S4) zones of the heart following MI. These data recapitulate those attained in  $\beta$ 2ARKO BMT mice (Fig. 2*A*), where decreased infiltration of monocytes/macrophages (Fig. 2*B*) and neutrophils (Fig. 2*D*) into the border and infarct (Fig. S4) zones were observed 4 d following MI. Interestingly, unlike in the  $\beta$ 2ARKO BMT mice, CCR2 inhibition did not impact mast cell infiltration (tryptase<sup>+</sup> cells; Fig. 2*A* and *C* and Fig. S2*E* and *G*). Further, post-MI survival, infarct size, and contractility did not differ between WT BMT and CCR2KO BMT mice (Fig. S5*A* and Table S2), although CCR2KO BMT mice had slightly less dilation following MI than WT BMT mice. These results suggest that altered CCR2 expression may be a major contributing, but not sole, factor to the decreased leukocyte recruitment response to the injured heart in mice lacking immune cell-expressed  $\beta$ 2AR.


**$\beta$ 2AR Stimulation Alters CCR2 Expression in a  $\beta$ -Arrestin2-Dependent Manner.** Because  $\beta$ 2ARKO BM has decreased expression of CCR2 compared with WT, we next sought to determine whether pharmacological activation of  $\beta$ 2AR reciprocally increases CCR2 expression. Thus, BM was isolated from WT C57BL/6 mice and

treated with the  $\beta$ 2AR-selective agonist salbutamol (Sal). Expression of CCR2 was quantified by RT-qPCR following Sal treatment over time. CCR2 levels were increased 6 and 24 h following  $\beta$ 2AR activation, demonstrating the ability to pharmacologically alter CCR2 levels using  $\beta$ 2AR ligands (Fig. 3*A*). Salbutamol-induced CCR2 expression observed in WT BM was not observed in  $\beta$ 2ARKO BM (Fig. 3*B*), confirming the specificity of the response. Migration assays were performed to determine whether  $\beta$ 2AR-dependent increases in CCR2 expression result in an enhanced functional response to CCL2-mediated migration. Indeed, CCL2-induced migration of WT BM was augmented with Sal pretreatment (Fig. 3*C* and *D*), whereas  $\beta$ 2ARKO BM did not migrate in response to CCL2, and Sal treatment had no effect on this response (Fig. 3*E* and *F*).

Because  $\beta$ 2AR stimulation engages both G protein- and  $\beta$ -arrestin ( $\beta$ ARR)-dependent signaling cascades, either of which may regulate downstream gene expression (8), we next sought to determine the proximal mechanism through which  $\beta$ 2AR stimulation increases CCR2 expression. Thus,  $\beta$ 2ARKO BM was infected with lentiviral constructs encoding either WT  $\beta$ 2AR (versus a GFP control



**Fig. 1. Effects of  $\beta$ 2ARKO on BM migration in response to chemokines.** (A) Representative Hoechst staining (white) from a 4-h migration assay of WT BM,  $\beta$ 2ARKO BM,  $\beta$ 2ARKO BM+lenti-GFP, and  $\beta$ 2ARKO BM+lenti- $\beta$ 2AR in response to CCL2 (100 ng/mL), CCL3 (100 ng/mL), or CXCL12 (10 ng/mL). A 1-h pretreatment with a CCR2 antagonist (10 nM) was used to inhibit CCR2-mediated migration. (B) Quantification of migration assay results. Values are expressed as fold over vehicle-stimulated migration.  $n = 4-8$ ; one-way ANOVA, \* $P < 0.05$  vs. WT BMT. (C) RT-qPCR was used to measure  $\beta$ 2AR and CCR2 expression in WT and  $\beta$ 2ARKO BM and  $\beta$ 2ARKO BM that had been transduced with either a GFP or  $\beta$ 2AR lentivirus.  $n = 4-8$ ; one-way ANOVA, \* $P < 0.05$  vs. WT BMT. Data are expressed as mean  $\pm$  SEM. NS, nonstimulated; RQ, relative quantitation.




**Fig. 2.** CCR2KO BMT reduces leukocyte infiltration into the heart following MI. (A) Representative CD68, trypsinase, and MPO staining for the border zone of hearts from WT C57BL/6 mice receiving WT, β2ARKO, or CCR2KO BMT that underwent MI surgery. Arrowheads indicate positive staining. *Insets* show higher magnification at 250 $\times$ . (B–D) Quantification of CD68 (B), trypsinase (C), and MPO (D) staining for the border zone of 4-d post-MI hearts from WT, β2ARKO, and CCR2KO BMT mice.  $n = 4$ –8; one-way ANOVA,  $^{\dagger}P < 0.01$  vs. WT BMT,  $^{*}P < 0.001$  vs. WT BMT; ns, not significant. Data are expressed as mean  $\pm$  SEM.

lentivirus), β2AR<sup>TY</sup> [lacking stimulatory G alpha subunit ( $G_{\alpha_s}$ ) coupling (9)], or β2AR<sup>GRK</sup> [deficient in G protein-coupled receptor kinase (GRK)-mediated phosphorylation and βARR recruitment (10)], and CCR2 expression was measured. Each lentiviral construct induced β2AR expression in β2ARKO BM to levels similar to WT BM, whereas GFP had no effect (Fig. S6A). Flag and GFP expression was also assessed by immunoblot to confirm transgene expression (Fig. S6B). WT β2AR and β2AR<sup>TY</sup> restored CCR2 expression in β2ARKO BM, whereas neither GFP nor β2AR<sup>GRK</sup> altered CCR2 expression (Fig. 4A). Functionally, migration in response to vehicle was unchanged by expression of any β2AR construct (Fig. 4B and C). However, corresponding to changes in CCR2 expression, β2AR<sup>GRK</sup> did not alter CCL2-mediated migration, whereas both WT β2AR and β2AR<sup>TY</sup> had enhanced migration in response to CCL2 (Fig. 4D and F). These results indicate that β2AR-mediated changes in CCR2 are dependent proximally upon βARR-dependent signaling. To confirm these results, BM was isolated from βARR1KO and βARR2KO BM, and CCR2 expression and migration responses were examined following treatment with Sal. As was observed in WT BM, Sal treatment increased CCR2 expression in βARR1KO BM (Fig. 5A) and resulted in enhanced migration in response to CCL2 (Fig. 5B and C). Conversely, Sal treatment of βARR2KO BM was unable to increase CCR2 expression (Fig. 5A) or CCL2-mediated migration (Fig. 5D and E). Thus, βARR2-dependent β2AR signaling increases CCR2 expression, thereby enhancing immune cell responsiveness to CCL2-mediated migration.

To further define the mechanism through which β2AR regulates CCR2 expression, transcription factor activation was examined using EMSAs. We assessed DNA binding of transcription factors reported to have putative binding sites in the CCR2 promoter and/or to regulate CCR2 expression [AP-1 (11), nuclear factor kappa-light-chain-enhancer of activated B cells (NF-κB) (12), and nuclear factor of activated T cells (NFAT) (13)], as well as cAMP response element binding protein (CREB) as a positive control, because it is known to be regulated downstream of βAR but with minimal impact on CCR2 transcription (14–16). AP-1 (Fig. 6A) and CREB (Fig. S7A) transcription factor binding were both decreased in BM from β2ARKO mice when compared with WT BM, whereas NF-κB (Fig. S7B) and NFAT (Fig. S7C) binding were unaltered between groups. We subsequently tested whether rescue of β2AR expression would restore transcription factor


binding in the β2ARKO BM and, as expected, canonical βARR-sensitive CREB DNA binding was restored upon reexpression of β2AR (Fig. S7D). Similarly, whereas GFP-infected β2ARKO BM still displayed reduced AP-1 DNA binding similar to β2ARKO alone (Fig. 6B), reexpression of WT β2AR restored AP-1 binding to WT levels. To determine the importance of AP-1 activation in the induction of CCR2 transcription, WT BM was pretreated with the AP-1 inhibitor SR11302 before Sal treatment. Increased CCR2 expression in response to Sal treatment was blocked by pretreatment with SR11302 (Fig. 6C). Human monocytes were also treated with Sal  $\pm$  SR11302, yielding identical results to those attained in mouse BM cells, highlighting the potential human relevance of our findings (Fig. 6D).

To determine whether βARR-dependent β2AR signaling is involved in the control of AP-1-dependent CCR2 expression, as suggested from our results above, β2ARKO BM was infected with the WT β2AR, β2AR<sup>TY</sup>, and β2AR<sup>GRK</sup> lentiviral constructs, or GFP control, and CCR2 expression was assessed. With restoration of WT β2AR expression, Sal increased CCR2,



**Fig. 3.** β2AR stimulation increases CCR2 expression and migration. (A) RT-qPCR was used to measure CCR2 expression in WT BM treated with vehicle (Veh) control or 1  $\mu$ M Sal over time (3 to 24 h).  $n = 3$ –8; one-way ANOVA,  $^{\dagger}P < 0.01$ ,  $^{*}P < 0.001$  vs. Veh. (B) RT-qPCR was used to measure CCR2 expression in WT and β2ARKO BM treated with Sal.  $n = 6$ ; one-way ANOVA,  $^{*}P < 0.001$  vs. Veh. (C) Representative Hoechst staining (white) from a 4-h migration assay of WT BM pretreated with vehicle or Sal and allowed to migrate in response to CCL2 (100 ng/mL). (D) Quantification of WT BM migration assay results. Values are expressed as fold over WT vehicle-stimulated migration.  $n = 7$ –8; one-way ANOVA,  $^{*}P < 0.05$ . (E) Representative Hoechst staining (white) from a 4-h migration assay of β2ARKO BM pretreated with vehicle or Sal and allowed to migrate in response to CCL2 (100 ng/mL). (F) Quantification of β2ARKO BM migration assay results. Values are expressed as fold over WT vehicle-stimulated migration.  $n = 7$ –8. Data are expressed as mean  $\pm$  SEM.





**Fig. 4.** Restoration of  $\beta 2AR$  expression reverses impairments in migration through  $\beta ARR$ -dependent mechanisms. (A) RT-qPCR was used to measure CCR2 expression in BM from WT or  $\beta 2ARKO$  BM transduced with GFP,  $\beta 2AR$ ,  $\beta 2AR^{TYY}$ , or  $\beta 2AR^{GRK-/-}$  lentivirus.  $n = 4-8$ ; one-way ANOVA,  $*P < 0.05$  vs. WT. (B) Representative Hoechst staining (white) from a 4-h migration assay in response to vehicle or CCL2 for  $\beta 2ARKO$  BM transduced with lentiviral constructs for GFP,  $\beta 2AR$ ,  $\beta 2AR^{TYY}$ , or  $\beta 2AR^{GRK-/-}$ . (C and D) Quantification of migration assay results. Values are expressed as fold over  $\beta 2ARKO + \beta 2AR$  Veh.  $n = 4-8$ ; one-way ANOVA,  $*P < 0.05$ ,  $^{\dagger}P < 0.01$ ,  $^{\ddagger}P < 0.001$ . Data are expressed as mean  $\pm$  SEM.

as previously seen in WT BM, which could be prevented with SR11302 pretreatment (Fig. 6E). GFP-infected cells were unresponsive to Sal  $\pm$  SR11302 (Fig. S7E). Similar to the WT  $\beta 2AR$  rescue,  $\beta 2AR^{TYY}$ -infected  $\beta 2ARKO$  BM had increased CCR2 expression following Sal treatment, which was blocked by SR11302 (Fig. 6F); however,  $\beta 2AR^{GRK-/-}$ -infected  $\beta 2ARKO$  BM showed no alteration in CCR2 expression with either Sal or SR11302 (Fig. S7F). These findings confirm that  $\beta ARR2$ -dependent  $\beta 2AR$  signaling controls CCR2 expression via regulation of AP-1 in hematopoietic cells.


**Restoration of  $\beta 2AR$ -Mediated  $\beta ARR$  Signaling in Vivo Reverses Leukocyte Dysfunction Following MI.** To determine whether restoration of  $\beta 2AR$  expression, in particular  $\beta 2AR$ -mediated  $\beta ARR$  signaling, in  $\beta 2ARKO$  BM could rescue the impaired cardiac leukocyte recruitment observed in post-MI  $\beta 2AR$  BMT mice,  $\beta 2ARKO$  BM was transduced with either the WT  $\beta 2AR$ ,  $\beta 2AR^{TYY}$ ,  $\beta 2AR^{GRK-/-}$ , or GFP control lentiviral constructs before BMT.  $\beta 2AR$  transcript expression in  $\beta 2ARKO$  BM following reconstitution was approximately that of endogenous levels for WT and mutant  $\beta 2AR$  (Fig. S6C), which corresponded to restoration of  $\beta AR$  membrane expression (Fig. S6D). Flag and GFP expression was assessed by immunoblot to confirm transgene expression in reconstituted BM (Fig. S6E). CCR2 expression was restored with the reexpression of WT  $\beta 2AR$  as well as with  $\beta 2AR^{TYY}$ , but not in either GFP- or  $\beta 2AR^{GRK-/-}$ -transduced BM (Fig. 7A). Analysis of the infarct size of animals showed similar surgical conditions between groups (Fig. S8). Leukocyte levels in the remote region (Fig. S9) were unchanged following MI; however, correlating with restoration of CCR2 expression, immunohistochemistry for monocyte/macrophages, mast cell and neutrophil infiltration to the border zone (Fig. 7B-E), and infarct (Fig. S10) of the injured myocardium of WT  $\beta 2AR$  and

$\beta 2AR^{TYY}$  were rescued compared with those of  $\beta 2ARKO + GFP$  and  $\beta 2AR^{GRK-/-}$  BMT mice.


## Discussion

Inflammatory processes are activated following acute cardiac injury, including chemokine-induced recruitment of immune cells to the site of injury, which are essential to mounting a reparative response and allowing subsequent healing (1). Sympathetic activity is known to regulate inflammation, with  $\beta 2AR$  being the predominant adrenergic receptor subtype involved in immunomodulation (4-6). Recent findings from our laboratory have identified a critical role for hematopoietically expressed  $\beta 2AR$  in survival following MI, wherein a lack of  $\beta 2AR$  on immune cells resulted in decreased leukocyte infiltration to the heart, failed scar formation, and cardiac rupture (7). Although splenic retention of leukocytes played a role in the phenotype, whether the diminished leukocyte recruitment to the injured heart in the absence of immune cell-expressed  $\beta 2AR$  involved an alteration in the response to promigratory chemokines was not determined. Thus, the purpose of this study was to determine whether immune cell-expressed  $\beta 2AR$  plays a key role in regulating chemokine responsiveness and leukocyte infiltration following acute cardiac injury.

Because trafficking of immune cells to sites of inflammation occurs through various chemokine receptors (2), we initially surveyed the expression levels of a number of the receptors in WT versus  $\beta 2ARKO$  BM, finding that CCR2 and CXCR4 levels in particular were significantly decreased. However, only CCR2 deficiency had a negative effect on BM cell migration in response to its ligand CCL2. This was in the absence of deficiencies in egress of cells from BM (7) that occur with global depletion of CCR2 (17, 18). Although the importance of CCL2 as a major chemoattractant of mononuclear cells to the ischemic heart has been extensively studied, the results are often in conflict, with both administration and inhibition of CCL2 showing improvements and detrimental effects in the remodeling following MI (19, 20). These discrepancies may be a result of the timing of administration or inhibition, because short-term elevations in CCL2 are protective whereas sustained elevations in CCL2 contribute to an enhanced progression toward heart failure (21-24). Indeed, studies examining the involvement of CCR2 following cardiac injury have shown beneficial effects with CCR2 inhibition (25-27), wherein both global and monocyte-directed RNAi-mediated deletion of CCR2 in mice that underwent MI surgery resulted in improved left ventricular remodeling (25, 26).



**Fig. 5.**  $\beta 2AR$  stimulation increases CCR2 expression and migration through  $\beta ARR2$ . (A) RT-qPCR was used to measure CCR2 expression in  $\beta ARR1KO$  and  $\beta ARR2KO$  treated with vehicle control or Sal. One-way ANOVA,  $*P < 0.05$  vs. Veh. (B-E) Representative Hoechst staining (white) and quantified results from a 4-h migration assay of  $\beta ARR1KO$  (B and C) and  $\beta ARR2KO$  BM (D and E) pretreated with Veh or Sal (1  $\mu M$ ; 24 h) before migration in response to Veh or CCL2 (100 ng/mL). Values are expressed as fold over Veh.  $n = 4-8$ ; one-way ANOVA,  $*P < 0.05$ . Data are expressed as mean  $\pm$  SEM.




**Fig. 6.** AP-1 activation is decreased in  $\beta 2$ ARKO BM. (A) Representative EMSAs from nuclear WT and  $\beta 2$ ARKO BM extracts incubated with an AP-1-specific probe. (B) A representative EMSA from nuclear WT and  $\beta 2$ ARKO BM extracts or  $\beta 2$ ARKO BM infected with lentiviral-expressed GFP or  $\beta 2$ AR and incubated with an AP-1-specific probe. (C) CCR2 expression was quantified in WT BM treated with vehicle, Sal  $\pm$  SR11302, or SR11302 alone.  $n = 4$ –8; one-way ANOVA,  $^{\dagger}P < 0.001$ . (D) Human monocytes were treated with vehicle, Sal  $\pm$  SR11302, or SR11302 alone. CCR2 expression was measured using RT-qPCR.  $n = 6$ –12; one-way ANOVA,  $^*P < 0.05$ ,  $^{\dagger}P < 0.01$ . (E and F)  $\beta 2$ ARKO BM was infected with lentiviral-expressed WT  $\beta 2$ AR (E) or lenti- $\beta 2$ AR<sup>TYY</sup> (F) and treated with vehicle, Sal  $\pm$  SR11302, or SR11302 alone. CCR2 expression was examined by RT-qPCR.  $n = 4$ –6; one-way ANOVA,  $^*P < 0.05$ ,  $^{\dagger}P < 0.01$ . Data are expressed as mean  $\pm$  SEM.

CCR2 is highly expressed on proinflammatory monocyte populations, and although  $\beta 2$ AR has been shown to modulate chemotaxis, our findings are novel in that they directly link decreased hematopoietic cell  $\beta 2$ AR expression with a corresponding reduction in CCR2 levels and CCL2-dependent migration. Previous studies have shown either no effect of sympathetic nervous system (SNS) stimulation on CCR2 expression in peripheral blood mononuclear cells (28) or decreased CCR2 expression in BM with SNS stimulation (29). These differences may be due to the lack of

specificity of norepinephrine and epinephrine treatment and activation of multiple adrenergic receptor subtypes. Further,  $\beta 2$ AR stimulation was previously shown in THP-1 human monocytic cells to increase CCR2 expression and migration through an undefined mechanism (30), whereas another study demonstrated that treatment of THP-1 cells with cAMP-elevating agents, including PDE3 inhibitors and dibutyl cAMP, decreased both CCR2 expression and CCL2-mediated migration (31). These results may suggest that  $G_{\alpha_s}$  protein-dependent  $\beta 2$ AR signaling acts to repress CCR2 expression.

Consistent with these reports, using  $\beta 2$ AR mutants lacking the ability to either engage  $G_{\alpha_s}$  protein signaling ( $\beta 2$ AR<sup>TYY</sup>) or to be phosphorylated by GRK ( $\beta 2$ AR<sup>GRK</sup>), we identified GRK-dependent  $\beta 2$ AR signaling as the mechanism by which  $\beta 2$ AR stimulation enhances CCR2 expression. Additionally, using BM cells from  $\beta$ ARRKO mice, we demonstrated that  $\beta$ ARR2, but not  $\beta$ ARR1, is specifically required for this effect.  $\beta$ ARR2 has been shown to regulate inflammatory responses in MI (32), as  $\beta$ ARR2KO mice had decreased survival after MI and decreased infiltration of macrophages to the infarct following MI, similar to our findings.  $\beta$ ARRs are known to be involved, either directly or indirectly, in the regulation of a number of transcription factors that could regulate CCR2 gene transcription, including AP-1 and NF- $\kappa$ B (33–37). Although the role of AP-1 has been minimally studied following MI (38, 39), it plays a well-established role in inflammation (40). We have demonstrated that  $\beta$ ARR2-dependent  $\beta 2$ AR signaling via AP-1 is required for CCR2 up-regulation in response to sympathetic stimulation, and that restoration of GRK/ $\beta$ ARR-dependent  $\beta 2$ AR signaling in immune cells rescues CCR2 expression, migration in response to CCL2, and cardiac infiltration following MI in vivo.

Although we show in our study that CCR2 deficiency in immune cells reduces their capacity for chemotaxis and infiltration to the heart post-MI, this deficiency was limited to the monocyte/macrophage and neutrophil populations in vivo, whereas mast cell infiltration was unchanged. This is in contrast to our previously reported study in which  $\beta 2$ ARKO chimeric mice had impairment in cardiac recruitment of all three cell types (7).



**Fig. 7.** Restoration of  $\beta 2$ AR expression in  $\beta 2$ ARKO BM reverses immune dysfunction following MI. (A) RT-qPCR was used to quantify CCR2 expression in reconstituted BM from mice that had GFP, WT  $\beta 2$ AR,  $\beta 2$ AR<sup>TYY</sup>, or  $\beta 2$ AR<sup>GRK</sup> transduced into  $\beta 2$ ARKO BM by lentivirus before transplantation. Values are expressed relative to WT BMT.  $n = 4$ –8; one-way ANOVA,  $^*P < 0.05$ ,  $^{\dagger}P < 0.001$  vs.  $\beta 2$ AR+GFP. (B) Representative CD68, tryptase, and MPO staining for the border zone of hearts from WT C57BL/6 mice receiving  $\beta 2$ ARKO BM with GFP, WT  $\beta 2$ AR,  $\beta 2$ AR<sup>TYY</sup>, or  $\beta 2$ AR<sup>GRK</sup> transduced before transplantation that underwent MI surgery. Arrowheads indicate positive staining. Insets show higher magnification at 250 $\times$ . (C–E) Quantification of CD68 (C), tryptase (D), and MPO (E) staining for the border zone of 4-d post-MI hearts from GFP, WT  $\beta 2$ AR,  $\beta 2$ AR<sup>TYY</sup>, or  $\beta 2$ AR<sup>GRK</sup> BMT mice. Arrowheads indicate positive staining.  $n = 4$ –8; one-way ANOVA,  $^*P < 0.05$ ,  $^{\dagger}P < 0.001$  vs.  $\beta 2$ ARKO+ $\beta 2$ AR BMT. Data are expressed as mean  $\pm$  SEM.



Further, in our current study, we did not observe an altered progression toward heart failure in CCR2KO BMT mice, which, coupled with the positive outcome of CCR2 inhibition in the aforementioned studies (25, 26) versus the catastrophic impact of immune cell-specific  $\beta$ 2AR deletion on cardiac remodeling following MI we previously reported (7), suggests that  $\beta$ 2ARKO chimeric mice likely have additional factors that contribute to their observed post-MI phenotype. For instance, we previously showed that enhanced vascular cell adhesion molecule 1 expression was associated with splenic retention of leukocytes in  $\beta$ 2ARKO BMT mice and that splenectomy partially restored cardiac leukocyte infiltration following MI (7), demonstrating that even with diminished CCR2 expression and responsiveness,  $\beta$ 2AR-deficient leukocytes retain some capacity to traffic to sites of injury in vivo. Thus, it may be beneficial to further investigate the role of CXCR4 expression, as well as a larger number of chemokine receptors and secreted factors from individual immune cell populations, to more fully elucidate how  $\beta$ 2AR controls these processes. The potential existence of multiple  $\beta$ 2AR-dependent mechanisms that could influence distinct immune cell populations suggests a widespread impact of  $\beta$ 2AR modulation on the regulation of early immune responses that could be targeted to alter post-MI recovery.

In summary, we have identified a role for  $\beta$ 2AR in the regulation of immune cell-specific CCR2 expression, where a lack of  $\beta$ 2AR expression in leukocytes results in decreased CCR2 expression, impaired migration to CCL2 in vitro, and decreased monocyte/macrophage and neutrophil cardiac infiltration following MI in vivo. Lentiviral-mediated reexpression of  $\beta$ 2AR in

$\beta$ 2ARKO BM before transplantation restored CCR2 expression and BM migration through a  $\beta$ ARR2-dependent pathway. These results demonstrate an immunomodulatory role for  $\beta$ ARR-biased  $\beta$ 2AR signaling in early immune responses following MI, which could be targeted to promote reparative processes while preventing chronic inflammatory events that are detrimental to healing. Further, because  $\beta$ -blockers are commonly used in patients around the time of acute cardiac ischemia, our results suggest they could impact the leukocyte-mediated repair response, warranting further investigation.

## Materials and Methods

**Surgery and Assays.** Detailed descriptions of coronary artery occlusion surgery, echocardiography, human monocyte cell culture, reverse transcription-quantitative PCR, migration assay, histological analysis, radioligand binding, immunoblot analysis, bone marrow transplant, bone marrow isolation, and lentiviral infection are provided in *SI Materials and Methods*. All animal procedures were performed in accordance with the guidelines of the Institutional Animal Care and Use Committee at the Temple University School of Medicine.

**Statistical Analysis.** Data presented are expressed as mean  $\pm$  SEM. Statistical analysis was performed using unpaired Student *t* tests, one-way ANOVA with a Tukey's multiple comparison test, or two-way repeated-measures ANOVA where appropriate using Prism 5.0 software (GraphPad Software), with *P* values indicated in the figure legends.

**ACKNOWLEDGMENTS.** This work was supported by NIH Grants HL105414 (to D.G.T.), HL091799 and HL085503 (to W.J.K.), and HL091804 (to C.J.T.) and an American Heart Association postdoctoral fellowship (to L.A.G.).

- Frangogiannis NG (2014) The inflammatory response in myocardial injury, repair, and remodelling. *Nat Rev Cardiol* 11(5):255–265.
- Turner MD, Nedjai B, Hurst T, Pennington DJ (2014) Cytokines and chemokines: At the crossroads of cell signalling and inflammatory disease. *Biochim Biophys Acta* 1843(11):2563–2582.
- Cavallera M, Frangogiannis NG (2014) Targeting the chemokines in cardiac repair. *Curr Pharm Des* 20(12):1971–1979.
- Nance DM, Sanders VM (2007) Autonomic innervation and regulation of the immune system (1987–2007). *Brain Behav Immun* 21(6):736–745.
- Elenkov IJ, Wilder RL, Chrousos GP, Vizi ES (2000) The sympathetic nerve—An integrative interface between two supersystems: The brain and the immune system. *Pharmacol Rev* 52(4):595–638.
- Padro CJ, Sanders VM (2014) Neuroendocrine regulation of inflammation. *Semin Immunol* 26(5):357–368.
- Grisanti LA, et al. (2016) Leukocyte-expressed  $\beta$ 2-adrenergic receptors are essential for survival after acute myocardial injury. *Circulation* 134(2):153–167.
- Lorton D, Bellinger DL (2015) Molecular mechanisms underlying  $\beta$ -adrenergic receptor-mediated cross-talk between sympathetic neurons and immune cells. *Int J Mol Sci* 16(3):5635–5665.
- Luttrell LM, et al. (2001) Activation and targeting of extracellular signal-regulated kinases by beta-arrestin scaffolds. *Proc Natl Acad Sci USA* 98(5):2449–2454.
- Shenoy SK, et al. (2006) Beta-arrestin-dependent, G protein-independent ERK1/2 activation by the beta2 adrenergic receptor. *J Biol Chem* 281(2):1261–1273.
- Phillips RJ, Lutz M, Premack B (2005) Differential signaling mechanisms regulate expression of CC chemokine receptor-2 during monocyte maturation. *J Inflamm (Lond)* 2:14.
- Li J, et al. (2001) Novel NEMO/I $\kappa$ B kinase and NF- $\kappa$ B target genes at the pre-B to immature B cell transition. *J Biol Chem* 276(21):18579–18590.
- Jung H, Miller RJ (2008) Activation of the nuclear factor of activated T-cells (NFAT) mediates upregulation of CCR2 chemokine receptors in dorsal root ganglion (DRG) neurons: A possible mechanism for activity-dependent transcription in DRG neurons in association with neuropathic pain. *Mol Cell Neurosci* 37(1):170–177.
- Tilley DG, Kim IM, Patel PA, Violin JD, Rockman HA (2009) Beta-arrestin mediates beta1-adrenergic receptor-epidermal growth factor receptor interaction and downstream signaling. *J Biol Chem* 284(30):20375–20386.
- Ueda A, et al. (1994) NF- $\kappa$ B and Sp1 regulate transcription of the human monocyte chemoattractant protein-1 gene. *J Immunol* 153(5):2052–2063.
- Yamamoto K, et al. (1999) Cloning and functional characterization of the 5'-flanking region of the human monocyte chemoattractant protein-1 receptor (CCR2) gene. Essential role of 5'-untranslated region in tissue-specific expression. *J Biol Chem* 274(8):4646–4654.
- Katayama Y, et al. (2006) Signals from the sympathetic nervous system regulate hematopoietic stem cell egress from bone marrow. *Cell* 124(2):407–421.
- Nakai A, Hayano Y, Furuta F, Noda M, Suzuki K (2014) Control of lymphocyte egress from lymph nodes through  $\beta$ 2-adrenergic receptors. *J Exp Med* 211(13):2583–2598.
- Ono K, et al. (1999) Prevention of myocardial reperfusion injury in rats by an antibody against monocyte chemoattractant and activating factor/monocyte chemoattractant protein-1. *Lab Invest* 79(2):195–203.
- Birdsall HH, et al. (1997) Complement C5a, TGF- $\beta$ 1, and MCP-1, in sequence, induce migration of monocytes into ischemic canine myocardium within the first one to five hours after reperfusion. *Circulation* 95(3):684–692.
- Wilson EM, Diwan A, Spinale FG, Mann DL (2004) Duality of innate stress responses in cardiac injury, repair, and remodeling. *J Mol Cell Cardiol* 37(4):801–811.
- Tarzami ST, Cheng R, Miao W, Kitsis RN, Berman JW (2002) Chemokine expression in myocardial ischemia: MIP-2 dependent MCP-1 expression protects cardiomyocytes from cell death. *J Mol Cell Cardiol* 34(2):209–221.
- Tarzami ST, et al. (2005) MCP-1/CCL2 protects cardiac myocytes from hypoxia-induced apoptosis by a G(alphai)-independent pathway. *Biochem Biophys Res Commun* 335(4):1008–1016.
- Dewald O, et al. (2005) CCL2/monocyte chemoattractant protein-1 regulates inflammatory responses critical to healing myocardial infarcts. *Circ Res* 96(8):881–889.
- Kaikita K, et al. (2004) Targeted deletion of CC chemokine receptor 2 attenuates left ventricular remodeling after experimental myocardial infarction. *Am J Pathol* 165(2):439–447.
- Majmudar MD, et al. (2013) Monocyte-directed RNAi targeting CCR2 improves infarct healing in atherosclerosis-prone mice. *Circulation* 127(20):2038–2046.
- Lavine KJ, et al. (2014) Distinct macrophage lineages contribute to disparate patterns of cardiac recovery and remodeling in the neonatal and adult heart. *Proc Natl Acad Sci USA* 111(45):16029–16034.
- Okutsu M, Suzuki K, Ishijima T, Peake J, Higuchi M (2008) The effects of acute exercise-induced cortisol on CCR2 expression on human monocytes. *Brain Behav Immun* 22(7):1066–1071.
- Xiu F, Stanojic M, Jeschke MG (2013) Norepinephrine inhibits macrophage migration by decreasing CCR2 expression. *PLoS One* 8(7):e69167.
- Guo YL, et al. (2014) Monocyte/macrophage  $\beta$ 2-AR as a target of antisymptomatic excitation-induced atherosclerotic progression. *Genet Mol Res* 13(4):8080–8088.
- Chuang SY, Yang SH, Pang JH (2011) Cilostazol reduces MCP-1-induced chemotaxis and adhesion of THP-1 monocytes by inhibiting CCR2 gene expression. *Biochem Biophys Res Commun* 411(2):402–408.
- Watarai K, Nakaya M, Nishida M, Kim KM, Kurose H (2013)  $\beta$ -Arrestin2 in infiltrated macrophages inhibits excessive inflammation after myocardial infarction. *PLoS One* 8(7):e68351.
- Witherow DS, Garrison TR, Miller WE, Lefkowitz RJ (2004) Beta-arrestin inhibits NF- $\kappa$ B activity by means of its interaction with the NF- $\kappa$ B inhibitor I $\kappa$ BaB. *Proc Natl Acad Sci USA* 101(23):8603–8607.
- Gao H, et al. (2004) Identification of beta-arrestin2 as a G protein-coupled receptor-stimulated regulator of NF- $\kappa$ B pathways. *Mol Cell* 14(3):303–317.
- Kang J, et al. (2005) A nuclear function of beta-arrestin1 in GPCR signaling: Regulation of histone acetylation and gene transcription. *Cell* 123(5):833–847.
- Kizaki T, et al. (2008) Beta2-adrenergic receptor regulates Toll-like receptor-4-induced nuclear factor- $\kappa$ B activation through beta-arrestin 2. *Immunology* 124(3):348–356.
- Wang Y, et al. (2006) Association of beta-arrestin and TRAF6 negatively regulates Toll-like receptor-interleukin 1 receptor signaling. *Nat Immunol* 7(2):139–147.
- Yoshida K, et al. (2001) Activation of mitogen-activated protein kinases in the non-ischemic myocardium of an acute myocardial infarction in rats. *Jpn Circ J* 65(9):808–814.
- Luo X, et al. (2009) c-Jun DNAs inhibit myocardial inflammation, ROS generation, infarct size, and improve cardiac function after ischemia-reperfusion injury. *Arterioscler Thromb Vasc Biol* 29(11):1836–1842.
- Schonthalder HB, Guinea-Viniegra J, Wagner EF (2011) Targeting inflammation by modulating the Jun/AP-1 pathway. *Ann Rheum Dis* 70(Suppl 1):i109–i112.